

MTH4103: Geometry I



Mid-term Test 2013

Duration: 40 minutes

Date and Time: 22nd February 2013, in the hour 13:00–14:00

Last name: _____

First name(s): _____

Student number: _____

The duration of this test is **40 minutes**. Answer **all** the questions. The marks (out of 100) allocated to each question are shown after the question numbers. For most questions, only the final answer to a question will be marked, so indicate this answer clearly. Simplify your answers as much as possible.

Calculators are **not** permitted in this examination. The unauthorised use of a calculator constitutes an examination offence.

Answer all questions in the spaces provided. You may do additional rough work on the backs of the question sheets or at the end, but this will not be looked at.

Do **not** turn over until an invigilator instructs you to do so.

For Examiner's use only

Question	Mark	Question	Mark
1		5	
2		6	
3		7	
4		8	
		Total (%)	

Question 1 (10 marks). Let $\mathbf{a} = \begin{pmatrix} -4 \\ 1 \\ -8 \end{pmatrix}$. Determine $|\mathbf{a}|$.

Question 2 (10 marks). Determine parametric equations for the line ℓ through the points $A = (-2, 1, 0)$ and $B = (4, 3, -2)$.

Question 3 (10 marks). Let $A = (-1, 1, 2)$ and $B = (-2, 3, 4)$, and let ℓ be the unique line on which A and B lie. Let P be the point on ℓ that is *not* between A and B and such that the distance from P to B is half the distance from A to B . (Thus $|\vec{AP}| = \frac{3}{2}|\vec{AB}|$.) Determine the position vector \mathbf{p} of P . [For notational convenience, A and B have position vectors \mathbf{a} and \mathbf{b} respectively.]

Question 4 (10 marks). Define carefully, but **without using coördinates**, the *scalar product* (or *dot product*) $\mathbf{u} \cdot \mathbf{v}$ of two vectors \mathbf{u} and \mathbf{v} .

Question 5 (20 marks). Apply Gaußian elimination to transform the following system of linear equations into echelon form (indicate this clearly), followed by back substitution to determine their full solution set. [You may assume that these are linear equations over \mathbb{R} in the variables x , y and z .]

$$\left. \begin{array}{l} y + 2z = -1 \\ x - 2y + 3z = -3 \\ -3x + 6y + 9z = -9 \end{array} \right\}.$$

Question 6 (20 marks). The four vertices of a regular tetrahedron of side length 2 can be taken to have position vectors $\mathbf{0}$, \mathbf{a} , \mathbf{b} and \mathbf{c} , where $\mathbf{a} = 2\mathbf{i}$, $\mathbf{b} = \mathbf{i} + \sqrt{3}\mathbf{j}$ and \mathbf{c} is dealt with below.

- (a) **(5 marks)** Calculate the vector product $\mathbf{a} \times \mathbf{b}$.
- (b) **(5 marks)** Let \mathbf{u} and \mathbf{v} be two vectors of length 2 at an angle of $\frac{\pi}{3}$ to each other. Calculate $\mathbf{u} \cdot \mathbf{v}$. [Hint: \mathbf{a} and \mathbf{b} are two such vectors.]
- (c) **(10 marks)** Given that \mathbf{c} has length 2 and has angle $\frac{\pi}{3}$ with both \mathbf{a} and \mathbf{b} , determine the two possibilities for \mathbf{c} .

Question 7 (10 marks). Exactly which of the following statements are true, and exactly which are false? [A statement of the form “ s_1 and s_2 are true” will *not* be taken to mean that you think s_3 , s_4 and s_5 are false.]

- (a) If \mathbf{u} , \mathbf{v} , \mathbf{w} are vectors such that \mathbf{u} is orthogonal to \mathbf{v} and \mathbf{v} is orthogonal to \mathbf{w} , then \mathbf{u} is orthogonal to \mathbf{w} .
- (b) If \mathbf{u} , \mathbf{v} , \mathbf{w} are coplanar then \mathbf{u} , \mathbf{v} , $\mathbf{u} \times \mathbf{v}$ are coplanar.
- (c) $\mathbf{v} \times \mathbf{u} = \mathbf{u} \times \mathbf{v}$ for all vectors \mathbf{u} and \mathbf{v} .
- (d) $\mathbf{v} \times \mathbf{u} \neq \mathbf{u} \times \mathbf{v}$ for all vectors \mathbf{u} and \mathbf{v} .
- (e) $\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w}) = \mathbf{w} \cdot (\mathbf{v} \times \mathbf{u})$ for all vectors \mathbf{u} , \mathbf{v} and \mathbf{w} .

Question 8 (10 marks). Let Π be the plane (in \mathbb{R}^3) with the equation $2x + y - 2z = 4$. Determine a point (called P) on Π and a nonzero vector (called \mathbf{n}) orthogonal to Π . Use this information to determine a point (called Q) at distance 6 from Π . [You may specify \mathbf{p} and \mathbf{q} instead of P and Q , where \mathbf{p} and \mathbf{q} are (respectively) the position vectors of P and Q .]

For rough work only.