

Microscopic Chaos, Fractals and Transport in Nonequilibrium Statistical Mechanics

Rainer Klages

Queen Mary University of London, School of Mathematical Sciences

Institut für Theoretische Elektrotechnik and
Institut für Theoretische Physik
Leibniz Universität Hannover, 30 May 2013



Outline

- 1 **Motivation:** microscopic chaos and transport; Brownian motion, dissipation and thermalization
- 2 the **thermostated dynamical systems approach** to nonequilibrium steady states and its surprising (fractal) properties
- 3 **generalized Hamiltonian dynamics** and universalities?

Why this talk?



W. and R. Mathis:
talks about **canonically
dissipative systems**




(sorry!)

but: R.F. Werner, *Generally observed features of the theory, like, e.g., the **approach of equilibrium** in macroscopic systems, deserve a general explanation don't they?*

main point of this talk: There is a **cross-link**...

Microscopic chaos in a glass of water?



- dispersion of a droplet of ink by *diffusion*
- assumption: *chaotic collisions* between billiard balls

microscopic chaos
↕
macroscopic transport

- relaxation to *equilibrium*

J.Ingenhousz (1785), R.Brown (1827), L.Boltzmann (1872),
P.Gaspard et al. (Nature, 1998)

Simple theory of Brownian motion

for a single **big tracer particle of velocity \mathbf{v}** immersed in a fluid:

$$\dot{\mathbf{v}} = -\kappa\mathbf{v} + \sqrt{\zeta}\xi(t) \quad \text{Langevin equation (1908)}$$


‘**Newton’s law of stochastic physics**’

force decomposed into

viscous damping

and

random kicks of surrounding particles



- models the interaction of a **subsystem** (tracer particle) with a **thermal reservoir** (fluid) in (\mathbf{r}, \mathbf{v}) -space
- two aspects: diffusion and dissipation; replace the tracer particle by a bottle of beer: **thermalization problem** in \mathbf{v} -space

Langevin dynamics

basic properties:

$$\dot{\mathbf{v}} = -\kappa\mathbf{v} + \sqrt{\zeta} \boldsymbol{\xi}(t)$$

stochastic

dissipative

not time reversible

⇒ **not Hamiltonian**

however:

see, e.g., **Zwanzig's (1973)** derivation of the Langevin equation from a heat bath of harmonic oscillators

non-Hamiltonian dynamics arises from **eliminating** the reservoir degrees of freedom by starting from a **purely Hamiltonian** system

Summary I

setting the scene:

- microscopic chaos and transport
- Brownian motion, dissipation and thermalization
- **Langevin dynamics: stochastic, dissipative, not time reversible, not Hamiltonian**

now to come:

the **deterministically thermostated dynamical systems approach** to nonequilibrium steady states

Nonequilibrium and the Gaussian thermostat

- Langevin equation with an electric field

$$\dot{\mathbf{v}} = \mathbf{E} - \kappa \mathbf{v} + \sqrt{\zeta} \boldsymbol{\xi}(t)$$

generates a **nonequilibrium steady state**: physical macro-scale quantities are **constant in time**

numerical inconvenience: slow relaxation

- alternative method via **velocity-dependent friction coefficient**

$$\dot{\mathbf{v}} = \mathbf{E} - \alpha(\mathbf{v}) \cdot \mathbf{v}$$

(for free flight); keep kinetic energy constant, $d\mathbf{v}^2/dt = 0$:


$$\alpha(\mathbf{v}) = \frac{\mathbf{E} \cdot \mathbf{v}}{v^2}$$

Gaussian (isokinetic) **thermostat**
Evans/Hoover (1983)

- follows from *Gauss' principle of least constraints*
- generates a *microcanonical velocity distribution*
- total *internal energy* can also be kept constant

The Lorentz Gas

free flight is a bit boring: consider the **periodic Lorentz gas** as a microscopic toy model for a conductor in an electric field




Galton (1877), Lorentz (1905)


couple it to a Gaussian thermostat - **surprise**: dynamics is **deterministic, chaotic, time reversible, dissipative, ergodic**

Hoover/Evans/Morriss/Posch (1983ff)

Gaussian dynamics: first basic property



Hoover, Moran (1989)



reversible equations of motion



fractal attractors in phase space



irreversible transport

Second basic property

- use equipartitioning of energy: $v^2/2 = T/2$

- consider ensemble averages: $\langle \alpha \rangle = \frac{E \cdot \langle \mathbf{v} \rangle}{T}$

absolute value of average **rate of phase space contraction**
= thermodynamic (Clausius) **entropy production**

that is:

entropy production is due to **contraction onto fractal attractor**
in nonequilibrium steady states

more generally: identity between Gibbs entropy production and
phase space contraction (Gerlich, 1973 and Andrey, 1985)

Third basic property

- define conductivity σ by $\langle \mathbf{v} \rangle =: \sigma \mathbf{E}$; into previous eq. yields

$$\sigma = \frac{T}{E^2} \langle \alpha \rangle$$

- combine with identity $-\langle \alpha \rangle = \lambda_+ + \lambda_-$ for Lyapunov exponents $\lambda_{+/-}$:

$$\sigma = -\frac{T}{E^2} (\lambda_+ + \lambda_-)$$

conductivity in terms of Lyapunov exponents


Posch, Hoover (1988); Evans et al. (1990)

similar relations for Hamiltonian dynamics and other transport coefficients from a different theory

Gaspard, Dorfman (1995)

Side remark: electrical conductivity

field-dependent electrical conductivity from NEMD computer simulations:



Lloyd et al. (1995)

- mathematical proof that there exists **Ohm's Law** for small enough (?) field strength (Chernov et al., 1993)
- but **irregular parameter dependence** of $\sigma(E)$ in simulations (cf. **book by RK, Part 1** on fractal transport coefficients)

Summary II

- **thermal reservoirs** needed to create steady states in nonequilibrium
- **Gaussian thermostat** as a deterministic alternative to Langevin dynamics
- Gaussian dynamics for **Lorentz gas** yields nonequilibrium steady states with very interesting dynamical properties

recall that Gaussian dynamics is *microcanonical*

last part:

construct a deterministic thermostat that generates a *canonical* distribution

The (dissipative) Liouville equation

Let $(\dot{\mathbf{r}}, \dot{\mathbf{v}})^* = \mathbf{F}(\mathbf{r}, \mathbf{v})$ be the equations of motion for a point particle and $\rho = \rho(t, \mathbf{r}, \mathbf{v})$ the probability density for the corresponding Gibbs ensemble

balance equation for **conserving the number of points** in phase space:

$$\frac{d\rho}{dt} + \rho \nabla \cdot \mathbf{F} = 0$$

Liouville equation (1838)

For Hamiltonian dynamics there is no phase space contraction, $\nabla \cdot \mathbf{F} = 0$, and **Liouville's theorem** is recovered:

$$\frac{d\rho}{dt} = 0$$

The Nosé-Hoover thermostat

Let $(\dot{\mathbf{r}}, \dot{\mathbf{v}}, \dot{\alpha})^* = \mathbf{F}(\mathbf{r}, \mathbf{v}, \alpha)$ with $\dot{\mathbf{r}} = \mathbf{v}$, $\dot{\mathbf{v}} = \mathbf{E} - \alpha(\mathbf{v})\mathbf{v}$ be the equations of motion for a point particle with **friction variable** α

problem: derive an equation for α that generates the **canonical distribution**

$$\rho(t, \mathbf{r}, \mathbf{v}, \alpha) \sim \exp \left[-\frac{v^2}{2T} - (\tau\alpha)^2 \right]$$

put the above equations into the Liouville equation

$$\frac{\partial \rho}{\partial t} + \dot{\mathbf{r}} \frac{\partial \rho}{\partial \mathbf{r}} + \dot{\mathbf{v}} \frac{\partial \rho}{\partial \mathbf{v}} + \dot{\alpha} \frac{\partial \rho}{\partial \alpha} + \rho \left[\frac{\partial \dot{\mathbf{r}}}{\partial \mathbf{r}} + \frac{\partial \dot{\mathbf{v}}}{\partial \mathbf{v}} + \frac{\partial \dot{\alpha}}{\partial \alpha} \right] = 0$$

restricting to $\partial \dot{\alpha} / \partial \alpha = 0$ yields the **Nosé-Hoover thermostat**

$$\dot{\alpha} = \frac{v^2 - 2T}{\tau^2 2T}$$

Nosé (1984), Hoover (1985)

widely used in NEMD computer simulations

Generalized Hamiltonian formalism for Nosé-Hoover

Dettmann, Morriss (1997): use the Hamiltonian

$$H(\mathbf{Q}, \mathbf{P}, Q_0, P_0) := e^{-Q_0} E(\mathbf{P}, P_0) + e^{Q_0} U(\mathbf{Q}, Q_0)$$

where $E(\mathbf{P}, P_0) = \mathbf{P}^2/(2m) + P_0^2/(2M)$ is the kinetic and $U(\mathbf{Q}, Q_0) = u(\mathbf{Q}) + 2TQ_0$ the potential energy of particle plus reservoir for **generalized** position and momentum coordinates

Hamilton's equations by imposing $H(\mathbf{Q}, \mathbf{P}, Q_0, P_0) = 0$:

$$\begin{aligned} \dot{\mathbf{Q}} &= e^{-Q_0} \frac{\mathbf{P}}{m}, \quad \dot{\mathbf{P}} = -e^{Q_0} \frac{\partial u}{\partial \mathbf{Q}} \\ \dot{Q}_0 &= e^{-Q_0} \frac{P_0}{M}, \quad \dot{P}_0 = 2(e^{-Q_0} E(\mathbf{P}, P_0) - e^{Q_0} T) \end{aligned}$$

uncoupled equations for $Q_0 = 0$ suggest **relation between physical and generalized coordinates**

$$\mathbf{Q} = \mathbf{q}, \quad \mathbf{P} = e^{Q_0} \mathbf{p}, \quad Q_0 = q_0, \quad P_0 = e^{Q_0} p_0$$

for $M = 2T\tau^2$, $\alpha = p_0/M$, $m = 1$ Nosé-Hoover recovered

note: the above transformation is **noncanonical!**

Nosé-Hoover dynamics

summary:

Nosé-Hoover thermostat constructed both from Liouville equation and from generalized Hamiltonian formalism

properties:

- fractal attractors
- identity between phase space contraction and entropy production
- formula for transport coefficients in terms of Lyapunov exponents

that is, we have the **same class as Gaussian dynamics**

basic question:

Are these properties **universal** for deterministic dynamical systems in nonequilibrium steady states altogether?

Non-ideal and boundary thermostats

counterexample 1:

increase the coupling for the Gaussian thermostat parallel to the field by making the friction **field-dependent**:

$$\dot{v}_x = E_x - \alpha(1 + E_x)v_x, \quad \dot{v}_y = -\alpha v_y$$

- **breaks the identity** between phase space contraction and entropy production and the conductivity-Lyapunov exponent formula
- **fractal attractors** seem to persist
- non-ideal Nosé-Hoover thermostat constructed analogously

counterexample 2:


a time-reversible deterministic boundary thermostat generalizing stochastic boundaries (**RK et al., 2000**)

- same results as above

Universality of Gaussian and Nosé-Hoover dynamics?

- ⊖ in general **no identity** between *phase space contraction and entropy production*
 - ⊖ consequently, relations between *transport coefficients and Lyapunov exponents* in thermostated systems are **not universal**
 - ⊕ existence of *fractal attractors* confirmed (stochastic reservoirs: open question)
- (possible way out: need to take a closer look at first problem...)

Outlook: the big picture




approach should be particularly useful for
small nonlinear systems

Acknowledgements and literature

counterexamples developed with:

K.Rateitschak (PhD thesis 2002, now Rostock), Chr.Wagner
(postdoc in Brussels 2002/3), G.Nicolis (Brussels)

literature:



(Part 2)