

# MAS115 Calculus I 2007-2008

Problem sheet for exercise class 8

- **Make sure you attend the exercise class that you have been assigned to!**
- The instructor will present the starred problems in class.
- You should then work on the other problems on your own.
- The instructor and helper will be available for questions.
- Solutions will be available online by Friday.

(\*) Problem 1:

[2007 exam questions]

Suppose that  $f$  has a negative derivative for all values of  $x$  and that  $f(1) = 0$ . Which of the following statements must be true of the function

$$h(x) = \int_0^x f(t) dt ?$$

- $h$  is a twice-differentiable function of  $x$ .
- $h$  and  $dh/dx$  are both continuous.
- The graph of  $h$  has a horizontal tangent at  $x = 1$ .
- $h$  has a local maximum at  $x = 1$ .
- $h$  has a local minimum at  $x = 1$ .
- The graph of  $h$  has an inflection point at  $x = 1$ .
- The graph of  $dh/dx$  crosses the  $x$ -axis at  $x = 1$ .

Problem 2: Sometimes it helps to reduce the integral step by step, using a trial substitution to simplify the integral a bit and then another to simplify it some more. Practice this on

$$\int \sqrt{1 + \sin^2(x-1)} \sin(x-1) \cos(x-1) dx .$$

- $u = x - 1$ , followed by  $v = \sin u$ , then by  $w = 1 + v^2$
- $u = \sin(x - 1)$ , followed by  $v = 1 + v^2$
- $u = 1 + \sin^2(x - 1)$

Problem 3: Suppose that  $f(x)$  is positive, continuous, and increasing over the interval  $[a, b]$ . By interpreting the graph of  $f$  show that

$$\int_a^b f(x) dx + \int_{f(a)}^{f(b)} f^{-1}(y) dy = bf(b) - af(a) .$$

Extra: Prove that

$$\int_0^x \left( \int_0^u f(t) dt \right) du = \int_0^x f(u)(x-u) du .$$

(Hint: Express the integral on the right hand side as the difference of two integrals. Then show that both sides of the equation have the same derivative with respect to  $x$ .)