

This is the author's version of a work that was accepted for publication in the Discrete Mathematics. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in *Disc. Math.* **290** (2005) 89–97.
<http://dx.doi.org/10.1016/j.disc.2004.10.006>

Multiple-elimination knockout tournaments with the fixed-win property

Matthew Fayers
Magdalene College, Cambridge, CB3 0AG, U.K.*

Abstract

We classify those triples (n, l, w) for which there exists a ‘knockout’ tournament for n players in which the winner always wins exactly w games and each loser loses exactly l games.

1 Introduction

In sporting competitions where a winner needs to be chosen in a relatively short time, a knockout tournament is frequently used. If more time is available, a double-elimination knockout tournament, in which a player or team is knocked out if it loses twice, may be employed. More generally, we may define an l -tuple-elimination knockout tournament to be one in which a player or team is eliminated if it loses l times. Specifically, an l -tuple elimination knockout tournament is a schedule in which:

- each game involves two players, one of whom wins and one of whom loses;
- the schedule for later games may depend upon the results of earlier games;
- if a player loses l games, he is eliminated from the tournament, and plays no further games;
- the tournament ends when exactly one player remains.

A great deal of literature is concerned with fairness in single- and (in [2]) double-elimination tournaments, assessing the probabilities of each player’s winning the tournament given his probabilities of beating each other player in a single game. Here, we approach the issue of fairness from a different angle, and consider the number of games a player needs to win in order to win the tournament. As far as we can tell, this issue has not been considered before. In the most frequently used double-elimination tournaments (such as in [1]), a player losing a game at an early stage will tend to need to win many more games in order to win the tournament than a player winning his early games. We say that an l -tuple elimination tournament for $n \geq 2$ players has the *fixed-win* property if, for some w , the winner of the tournament is guaranteed to have won exactly w games. We refer to such a tournament as a $FW(n, l, w)$.

The aim of this paper is to classify those triples (n, l, w) for which a $FW(n, l, w)$ exists. Two obvious examples are as follows.

*Current address: Queen Mary University of London, Mile End Road, London, E1 4NS, U.K.

- The standard single-elimination knockout tournament for 2^w players is a $\text{FW}(2^w, 1, w)$.
- Two players playing a ‘first to w ’ match is a $\text{FW}(2, w, w)$.

In fact, these are almost the only examples. Our main theorem is as follows.

Theorem 1.1. *Suppose that $n \geq 2$, and that l, w are positive integers. Then a $\text{FW}(n, l, w)$ exists if and only if*

- $l = 1$ and $n = 2^w$,
- $n = 2$ and $l = w$ or
- $(n, l, w) = (16, 2, 6)$.

2 The proof of Theorem 1.1

In order to prove Theorem 1.1, we need to prove a more general result. Given $n \geq 1$ and positive integers $w_1, \dots, w_n, l_1, \dots, l_n$, we ask whether we can arrange a tournament for n players (numbered $1, \dots, n$) in which $n - 1$ players are eliminated and the remaining player wins the tournament, and in which:

- player i is eliminated if and only if he loses l_i games;
- player i wins the tournament if and only if he wins w_i games.

We call such a tournament a $T(w_1, \dots, w_n, l_1, \dots, l_n)$, and say that the matrix

$$\begin{pmatrix} w_1 & \dots & w_n \\ l_1 & \dots & l_n \end{pmatrix}$$

is good if a $T(w_1, \dots, w_n, l_1, \dots, l_n)$ exists. In particular, a $\text{FW}(n, l, w)$ exists if and only if the $2 \times n$ matrix

$$\begin{pmatrix} w & \dots & w \\ l & \dots & l \end{pmatrix}$$

is good. We shall classify good matrices; clearly

$$\begin{pmatrix} w_1 & \dots & w_n \\ l_1 & \dots & l_n \end{pmatrix}$$

is good if and only if

$$\begin{pmatrix} w_{\sigma(1)} & \dots & w_{\sigma(n)} \\ l_{\sigma(1)} & \dots & l_{\sigma(n)} \end{pmatrix}$$

is good for some permutation σ .

Given a matrix

$$\mathbf{M} = \begin{pmatrix} w_1 & \dots & w_n \\ l_1 & \dots & l_n \end{pmatrix}$$

and given $1 \leq i < j \leq n$, we define the (i, j) -descendants of \mathbf{M} to be the matrices

$$\begin{pmatrix} w_1 & \dots & w_i - 1 & \dots & w_j & \dots & w_n \\ l_1 & \dots & l_i & \dots & l_j - 1 & \dots & l_n \end{pmatrix}$$

and

$$\begin{pmatrix} w_1 & \dots & w_i & \dots & w_j - 1 & \dots & w_n \\ l_1 & \dots & l_i - 1 & \dots & l_j & \dots & l_n \end{pmatrix}$$

where we delete any column in which the lower entry is 0.

Proposition 2.1. *The matrix $\begin{pmatrix} w_1 \\ l_1 \end{pmatrix}$ is good if and only if $w_1 = 0$. If $n \geq 2$, then the matrix $\begin{pmatrix} w_1 & \dots & w_n \\ l_1 & \dots & l_n \end{pmatrix}$ is good if and only if for some $1 \leq i < j \leq n$ both of the (i, j) -descendants are good.*

Proof. For $n = 1$, the result is obvious: since there is only one player to begin with, this player has won the tournament winning no games. If $n \geq 2$, suppose that we wish to construct a $T(w_1, \dots, w_n, l_1, \dots, l_n)$ in which the first game is between players i and j . If i beats j , then the remainder of the tournament may be viewed as a $T(w_1, \dots, w_{i-1}, w_i - 1, w_{i+1}, \dots, w_n, l_1, \dots, l_{j-1}, l_j - 1, l_{j+1}, \dots, l_n)$, while if j beats i , then the remainder of the tournament is a $T(w_1, \dots, w_{j-1}, w_j - 1, w_{j+1}, \dots, w_n, l_1, \dots, l_{i-1}, l_i - 1, l_{i+1}, \dots, l_n)$. So a $T(w_1, \dots, w_n, l_1, \dots, l_n)$ exists if and only if both the latter two tournaments exist, for some i and j . \square

Proposition 2.2. *The matrix $\begin{pmatrix} w_1 & w_2 \\ l_1 & l_2 \end{pmatrix}$ is good if and only if $w_1 = l_2$ and $l_1 = w_2$.*

Proof. This follows easily by induction and Proposition 2.1. \square

It will turn out that there are remarkably few good matrices. We begin with the case where each l_i equals 1.

Proposition 2.3. *The matrix*

$$\mathbf{M} = \begin{pmatrix} w_1 & \dots & w_n \\ 1 & \dots & 1 \end{pmatrix}$$

is good if and only if $\sum_{k=1}^n 2^{-w_k} = 1$.

To prove this, we need the following simple lemma.

Lemma 2.4. *If $n > 1$ and a_1, \dots, a_n are integers such that $2^{a_1} + \dots + 2^{a_n} = 2^a$ for some integer a , then $a_i = a_j$ for some $i \neq j$.*

Proof. If the a_i are all distinct, let a_m be the smallest. Then $2^{a_1} + \dots + 2^{a_n}$ is an odd integer multiple of 2^{a_m} , and so cannot be a power of 2 (since it does not equal 2^{a_m}). \square

Proof of Proposition 2.3. The case $n = 2$ follows from Proposition 2.2. For $n > 2$, suppose that \mathbf{M} is good, and let i and j be as in Proposition 2.1. Then by induction we must have

$$2^{-(w_i-1)} + \sum_{k \neq i, j} 2^{-w_k} = 1 \tag{1}$$

and

$$2^{-(w_j-1)} + \sum_{k \neq i, j} 2^{-w_k} = 1. \tag{2}$$

In particular, we have $w_i = w_j$, and so

$$\sum_{k=1}^n 2^{-w_k} = 1. \quad (3)$$

Conversely, suppose that $n \geq 3$ and that (3) holds. By Lemma 2.4 we can find i and j such that $w_i = w_j$; then equations (1–2) above hold, and so by induction both of the (i, j) -descendants of \mathbf{M} are good. \square

Now we consider the case where some of the l_i equal 2.

Proposition 2.5. *Suppose $n \geq 3$. Then the matrix*

$$\mathbf{M} = \begin{pmatrix} u_1 & \cdots & u_{n-1} & v \\ 1 & \cdots & 1 & 2 \end{pmatrix}$$

is good if and only if

$$v = 1 \text{ or } 2 \quad \text{and} \quad \sum_{k=1}^{n-1} 2^{-u_k} = \frac{v}{4}.$$

Proof. We begin with the case $n = 3$. The $(1, 2)$ -descendants of \mathbf{M} are

$$\begin{pmatrix} u_1 - 1 & v \\ 1 & 2 \end{pmatrix} \text{ and } \begin{pmatrix} u_2 - 1 & v \\ 1 & 2 \end{pmatrix},$$

and these are both good if and only if $v = 1$ and $u_1 = u_2 = 3$. The $(1, 3)$ -descendants are

$$\begin{pmatrix} u_1 - 1 & u_2 & v \\ 1 & 1 & 1 \end{pmatrix} \text{ and } \begin{pmatrix} u_2 & v - 1 \\ 1 & 2 \end{pmatrix},$$

and these are both good if and only if $u_2 = 2$, $v - 1 = 1$ and $2^{-(u_1-1)} + 2^{-u_2} + 2^{-v} = 1$, i.e. if and only if $(u_1, u_2, v) = (2, 2, 2)$; similarly for the $(2, 3)$ -descendants.

Now we assume $n > 3$, and suppose that the (i, j) -descendants of \mathbf{M} are both good. There are two cases to consider.

$[i, j < n]$ By induction we have

$$2^{-(u_i-1)} + \sum_{k \neq i, j} 2^{-u_k} = \frac{v}{4} = 2^{-(u_j-1)} + \sum_{k \neq i, j} 2^{-u_k},$$

and so $u_i = u_j$. Thus $\sum_{k=1}^{n-1} 2^{-u_k}$ equals $\frac{v}{4}$ as well, and so \mathbf{M} satisfies one of the above criteria.

$[i < j = n]$ Now the (i, j) -descendants of \mathbf{M} are

$$\begin{pmatrix} u_1 & \cdots & u_{i-1} & u_i - 1 & u_{i+1} & \cdots & u_{n-1} & v \\ 1 & \cdots & 1 & 1 & 1 & \cdots & 1 & 1 \end{pmatrix} \text{ and } \begin{pmatrix} u_1 & \cdots & u_{i-1} & u_{i+1} & \cdots & u_{n-1} & v - 1 \\ 1 & \cdots & 1 & 1 & \cdots & 1 & 2 \end{pmatrix},$$

so by induction we must have $v - 1 = 1$ or 2 and

$$\sum_{k \neq i} 2^{-u_k} = \frac{v-1}{4}, \quad \sum_{k \neq i} 2^{-u_k} + 2^{-(u_i-1)} + 2^{-v} = 1.$$

This gives

$$2^{-(u_i-1)} = 1 - 2^{-v} - \frac{v-1}{4};$$

if $v - 1 = 2$, then we get $2^{-(u_i-1)} = \frac{3}{8}$, which is impossible, so we have $v - 1 = 1$, which gives $2^{-(u_i-1)} = \frac{1}{2}$, whence $u_i = 2$ and we have $v = 2$, $\sum_{k=1}^{n-1} 2^{-u_k} = \frac{1}{2}$.

Conversely, suppose that \mathbf{M} satisfies the conditions of the proposition. By Lemma 2.4 we must have $u_i = u_j$ for some $1 \leq i < j < n$; then the (i, j) -descendants of \mathbf{M} both satisfy this criterion as well. \square

In order to deal with the case where two or more of the l_i equal 2, we need to introduce a certain function. Suppose S is a finite multiset of integers such that $\sum_{a \in S} 2^{-a} = \frac{1}{4}$. Then we define

$$h(S) = \frac{3}{4} - \sum_{a \in S} a 2^{-a-1}.$$

For example, we have

$$\begin{aligned} h(\{2\}) &= \frac{1}{2}, & h(\{3, 3\}) &= \frac{3}{8}, & h(\{3, 4, 4\}) &= \frac{5}{16}, \\ h(\{3, 4, 5, 5\}) &= \frac{9}{32}, & h(\{4, 4, 4, 4\}) &= \frac{1}{4}. \end{aligned}$$

Note that the function h satisfies

- $h(\{2\}) = \frac{1}{2}$, and
- $h(\{a_1, a_1, a_2, \dots, a_n\}) = h(\{a_1 - 1, a_2, \dots, a_n\}) - 2^{-a_1}$.

Since by Lemma 2.4 any finite multiset S consisting of at least two integers and satisfying $\sum_{a \in S} 2^{-a} = \frac{1}{4}$ must contain two equal elements, these rules are enough to compute h recursively.

A special case of h is the following.

Lemma 2.6. For $a \geq 2$, let S be the multiset with 2^{a-2} elements all equal to a . Then

$$h(S) = \frac{6-a}{8}.$$

Now we can deal with the general case in which each l_i equals 1 or 2.

Proposition 2.7. Suppose that $n \geq 3$ and $0 \leq r \leq n - 2$. Then the matrix

$$\mathbf{M} = \begin{pmatrix} u_1 & \dots & u_r & v_{r+1} & \dots & v_n \\ 1 & \dots & 1 & 2 & \dots & 2 \end{pmatrix}$$

is good if and only if

$$\sum_{k=r+1}^n 2^{-v_k} = \frac{1}{4} \quad \text{and} \quad \sum_{k=1}^r 2^{-u_k} = h(\{v_{r+1}, \dots, v_n\}).$$

Proof. For the ‘if’ part, we have by Lemma 2.4 that $v_i = v_j$ for some $i \neq j$. Then, by the recursive description of h and by induction (or by Proposition 2.5 in the case $r = n - 2$), the (i, j) -descendants of \mathbf{M} are both good.

For the ‘only if’ part, we proceed by induction on $n - r$ and r , beginning with the case $n = 3, r = 1$; notice that the conditions of the proposition cannot hold here. The $(1, 2)$ -descendants of

$$\begin{pmatrix} u_1 & v_2 & v_3 \\ 1 & 2 & 2 \end{pmatrix}$$

are

$$\begin{pmatrix} u_1 - 1 & v_2 & v_3 \\ 1 & 1 & 2 \end{pmatrix} \text{ and } \begin{pmatrix} v_2 - 1 & v_3 \\ 2 & 2 \end{pmatrix}.$$

For both of these to be good we should need $v_2 = 3, v_3 = 2$ and $2^{-(u_1-1)} + 2^{-v_2} = \frac{v_3}{4}$, which is impossible. Similarly for the $(1, 3)$ -descendants. The $(2, 3)$ -descendants are

$$\begin{pmatrix} u_1 & v_2 - 1 & v_3 \\ 1 & 2 & 1 \end{pmatrix} \text{ and } \begin{pmatrix} u_1 & v_2 & v_3 - 1 \\ 1 & 1 & 2 \end{pmatrix};$$

similarly, these cannot both be good.

Now we examine the case $r > 1, n - r = 2$, and we suppose that the (i, j) -descendants of \mathbf{M} are both good. We look at the various possibilities for i, j .

$[i, j \leq r]$ By induction we have $2^{-v_{n-1}} + 2^{-v_n} = \frac{1}{4}$ and

$$\sum_{k \neq i, j} 2^{-u_k} + 2^{-(u_i-1)} = h(\{v_{n-1}, v_n\}) = \sum_{k \neq i, j} 2^{-u_k} + 2^{-(u_j-1)},$$

so that $u_i = u_j$ and thus

$$\sum_{k=1}^r 2^{-u_k} = h(\{v_{n-1}, v_n\})$$

and \mathbf{M} satisfies the criteria of the proposition.

$[(i, j) = (n - 1, n)]$ By Proposition 2.5 we have

$$v_m - 1 = 1 \text{ or } 2, \quad \sum_{k=1}^r 2^{-u_k} + 2^{-v_{2n-1-m}} = \frac{v_m - 1}{4}$$

for $m = n - 1, n$. This gives a contradiction unless we have $v_{n-1} = v_n = 3$, whence $2^{-v_{n-1}} + 2^{-v_n} = \frac{1}{4}$ and $\sum_{k=1}^r 2^{-u_k} = \frac{3}{8}$, as required.

$[i \leq r < j]$ By induction we must have

$$v_j - 1 = v_{2n-1-j} = 3, \quad \sum_{k \neq i} 2^{-u_k} = \frac{3}{8}$$

while by Proposition 2.5 we need

$$v_{2n-1-j} = 1 \text{ or } 2,$$

a contradiction.

Finally, we examine the case where $n - r > 2$, and suppose that the (i, j) -descendants of \mathbf{M} are good.

$[i, j \leq r]$ Exactly as above we find that $u_i = u_j$ and that \mathbf{M} satisfies the criteria of the proposition.

$[i, j > r]$ By induction we have

$$\sum_{k \neq i, j} 2^{-v_k} + 2^{-(v_i-1)} = \frac{1}{4} = \sum_{k \neq i, j} 2^{-v_k} + 2^{-(v_j-1)}$$

so that $v_i = v_j$, and

$$\begin{aligned} \sum_{k=1}^r 2^{-u_k} + 2^{-v_i} &= h(v_{r+1}, \dots, v_{i-1}, v_{i+1}, \dots, v_{j-1}, v_{j+1}, \dots, v_n, v_j - 1) \\ &= h(v_{r+1}, \dots, v_n) + 2^{-v_j}, \end{aligned}$$

as required.

$[i \leq r < j]$ In this case we have

$$\sum_{k \neq j} 2^{-v_k} = \frac{1}{4} = \sum_{k \neq j} 2^{-v_k} + 2^{-(v_j-1)},$$

a contradiction. □

Now we consider those cases where one of the l_i is at least 3.

Proposition 2.8. *If $n \geq 2$ and $x \geq 3$, then the matrix*

$$\mathbf{M} = \begin{pmatrix} u_1 & \dots & u_{n-1} & v \\ 1 & \dots & 1 & x \end{pmatrix}$$

is good if and only if

$$v = 1 \quad \text{and} \quad \sum_{k=1}^{n-1} 2^{-u_k} = 2^{-x}.$$

Proof. We use induction on n and x , with the case $n = 2$ being a special case of Proposition 2.2. If $n > 2$ and \mathbf{M} satisfies the hypotheses, then $u_i = u_j$ for some $i \neq j$, and the (i, j) -descendants are good, by induction.

Now suppose that $n > 2$ and that the (i, j) -descendants of \mathbf{M} are both good. If $i, j < n$, then by induction we have $v = 1$ and

$$2^{-(u_i-1)} + \sum_{k \neq i, j} 2^{-u_k} = 2^{-x} = 2^{-(u_j-1)} + \sum_{k \neq i, j} 2^{-u_k},$$

so that $u_i = u_j$ and \mathbf{M} satisfies the hypotheses. If $i < j = n$, then by induction we have both $v - 1 = 1$ and $v = 1$ – contradiction – unless $x = 3$, when we may have

$$v = 2, \sum_{k \neq i} 2^{-u_k} = 2^{-x}, \sum_{k \neq i} 2^{-u_k} + 2^{-(u_i-1)} = \frac{1}{2};$$

but this gives $2^{-(u_i-1)} = \frac{3}{8}$, also a contradiction. □

Now we show that we have found all good matrices.

Proposition 2.9. *If $n \geq 3$ and $l_i \geq 3, l_j \geq 2$ for some $i \neq j$, then the matrix*

$$\mathbf{M} = \begin{pmatrix} w_1 & \dots & w_n \\ l_1 & \dots & l_n \end{pmatrix}$$

is not good.

Proof. Suppose that the (i, j) -descendants of \mathbf{M} are both good. Then by induction neither of the (i, j) -descendants can satisfy the conditions of the proposition, and so we must have $l_i = 3, l_j = 2$ and $l_k = 1$ for all $k \neq i, j$ (if $n = 3$ then are two extra possible cases, but these are easily dealt with). But then we require $w_i = w_j - 1 = 3$ by Proposition 2.7, while $w_i - 1 = 1$ by Proposition 2.8. Contradiction. \square

Proof of Theorem 1.1. As noted above, a $\text{FW}(n, l, w)$ exists if and only if the matrix

$$\mathbf{M} = \begin{pmatrix} w & \dots & w \\ l & \dots & l \end{pmatrix}$$

is good. For $n = 2$, this is true if and only if $l = w$, by Proposition 2.2. For $l = 1$, we need $n \cdot 2^{-w} = 1$, so that $n = 2^w$. If $n > 2$ and $l = 2$, then by Proposition 2.7 we need $n \cdot 2^{-w} = \frac{1}{4}$, and $h(w, \dots, w) = 0$. By Lemma 2.6, this happens if and only if $n = 16, w = 6$.

If $n, l > 2$, then \mathbf{M} is not good, by Proposition 2.9. \square

3 Tournaments with multiple winners

An interesting extension to this problem is to ask about tournaments which produce multiple winners; this is applicable to the situation where a set of players is chosen to proceed to the next stage of a competition. We define a $\text{FW}_r(n, l, w)$ to be a competition for n players which produces r winners, each of whom has won exactly w games, and $n - r$ losers, each of whom has lost exactly l games. Classifying such tournaments seems to be rather harder than in the case $r = 1$; the following simple observations provide lots of ‘trivial’ examples of these tournaments:

1. a $\text{FW}_r(n, l, w)$ exists if and only if a $\text{FW}_{n-r}(n, w, l)$ exists;
2. if a $\text{FW}_r(n, l, w)$ and a $\text{FW}_s(m, l, w)$ exist, then a $\text{FW}_{r+s}(n + m, l, w)$ exists.

But there are other examples. For instance, it is fairly easy to show that for any $r \geq 1$ a $\text{FW}_r(2^{8r-4}, 2, 8r - 2)$ exists. Of course, the case $r = 1$ gives the ‘sporadic’ case $(16, 2, 6)$ of Theorem 1.1. We hope to be able to say more about tournaments with multiple winners in a future paper.

References

- [1] C. Edwards, ‘Double-elimination tournaments: counting and calculating’, *Amer. Statist.* **50** (1996), no. 1, 27–33.
- [2] E. Jaryszak, P. Thompson & J. Wamil, ‘Reducing the pairing effect in 4-team double-elimination tournaments’, *Math. Sci.* **25** (2000), no. 2, 110–121.