

Some measures of finite groups related to permutation bases

Peter J. Cameron

School of Mathematical Sciences
Queen Mary, University of London
Mile End Road
London E1 4NS
U.K.

p.j.cameron@qmul.ac.uk

Abstract

I define three “measures” of the complicatedness of a finite group in terms of bases in permutation representations of the group, and consider their relationships to other measures.

1 Introduction and history

The purpose of this paper is to define and investigate three functions which in some sense “measure” the size or complicatedness of the group. The functions are defined in minimax fashion in terms of base size in permutation representations of the group. The values of these measures for symmetric groups will be given.

I begin with a brief account of some similar measures, which are related to the new measures in various ways.

If f is a function from (isomorphism classes of) finite groups to natural numbers, then f' is the function defined by

$$f'(G) = \max\{f(H) : H \leq G\}.$$

The function f' is necessarily monotonic: that is, if $H \leq G$, the $f'(H) \leq f'(G)$. Conversely, if f is monotonic, then $f = f'$; so every monotonic function is of this form.

The first such measure is $d(G)$, the minimum number of generators of G . Now it is well-known that $d(S_n) = 2$; so this invariant is not very sensitive.

However, d' is more interesting. Babai [1] raised the question of determining $d'(S_n)$. The motivation was computational group theory: if we are given an arbitrary (possibly very large) subset of S_n , and want to know the subgroup they generate, then we know that we can replace the given set by a set of size at most $d'(S_n)$ generating the same subgroup. This suggests that we also want, if possible, an algorithmic method for finding such a minimum-size generating set.

Babai considered the related function l , where $l(G)$ is the length of the longest chain of subgroups of G . Since, obviously, $d(G) \leq l(G)$, and $l(G)$ is monotonic (that is, $l(G) = l'(G)$), we see that $d'(G) \leq l(G)$ for any group G . Babai proved that

$$d'(S_n) \leq l(S_n) \leq 2n - 1.$$

The exact value of $l(S_n)$ was established by Cameron, Solomon and Turull [3]:

$$l(S_n) = \left\lceil \frac{3n}{2} \right\rceil - b(n) - 1,$$

where $b(n)$ is the number of 1s in the base 2 representation of n .

Jerrum [5] showed that $d'(S_n) \leq n - 1$. His proof was algorithmic; given an arbitrary set of permutations, a set of at most $n - 1$ permutations generating the same subgroup can be found efficiently.

The exact value of $d'(S_n)$ was established by McIver and Neumann [6]:

$$d'(S_n) = \left\lfloor \frac{n}{2} \right\rfloor \text{ for } n > 3.$$

Whiston [7] considered the invariant $\mu(G)$, the maximal size of an independent generating set for G , where a set is *independent* if none of its elements lies in the subgroup generated by the others. Since any independent set is an independent generating set for the subgroup it generates, $\mu'(G)$ is the maximum size of an independent subset of G . The parameter $\mu(G)$ appears in the work of Diaconis and Saloff-Coste [4] on the rate of convergence of the product replacement algorithm for finding a random element of a finite group.

Whiston showed that

$$\mu(S_n) = \mu'(S_n) = n - 1.$$

However, he observed that there are groups with $\mu'(G) > \mu(G)$.

Cameron and Cara [2] found all independent generating sets of size $n - 1$ in S_n .

2 The base measures

Let G be a permutation group on Ω . A *base* for G a sequence of points of Ω whose pointwise stabiliser is the identity. (Treating a base as a sequence rather than a set fits with the use of bases in computational group theory, where the elements of a base are chosen in order.) A base is called *irredundant* if no point is fixed by the pointwise stabiliser of its predecessors; it is *minimal* if no point is fixed by the pointwise stabiliser of the other points in the sequence.

It is computationally a simple matter to choose an irredundant base; simply choose each base point to be a point moved by the stabiliser of the points previously chosen. It is less straightforward to choose a minimal base. Note that a base is minimal if and only if every re-ordering of it is irredundant.

Now, given a finite group G , we define three numbers $b_1(G)$, $b_2(G)$, $b_3(G)$, as follows. In each case, the maximum is taken over all permutation representations of G (not necessarily faithful).

- $b_1(G)$ is the maximum, over all representations, of the maximum size of an irredundant base;
- $b_2(G)$ is the maximum, over all representations, of the maximum size of a minimal base;
- $b_3(G)$ is the maximum, over all representations, of the minimum base size.

Clearly we have:

Proposition 2.1 $b_3(G) \leq b_2(G) \leq b_1(G)$. □

These inequalities can be strict. The group $G = \text{PSL}(2, 7)$ has $b_1(G) = 5$, $b_2(G) = 4$, and $b_3(G) = 3$.

Now $b_1(G)$ is a parameter we have seen before!

Proposition 2.2 $b_1(G) = l(G)$.

Proof Given an irredundant base of size $b_1(G)$, the stabilisers of its initial sequences form a properly descending chain of subgroups of length $b_1(G)$. So $b_1(G) \leq l(G)$.

Conversely, let

$$G = G_0 > G_1 > \cdots > G_l = 1$$

be a chain of subgroups of length $l = l(G)$. Consider the action on the union of the coset spaces of the subgroups G_i , and let α_i be the point G_i of the coset space $(G : G_i)$. Then $(\alpha_1, \dots, \alpha_l)$ is an irredundant base. So $l(G) \leq b_1(G)$. □

We will see in the next section a connection between $b_2(G)$ and $\mu(G)$. I know much less about $b_3(G)$. One observation is the following:

Proposition 2.3 *Let G be a non-abelian finite simple group. Then $b_3(G)$ can be calculated by considering only the primitive permutation representations of G .*

Proof Given any permutation representation of G , we can discard fixed points, so that G acts faithfully on each orbit. Now let b_3^*G be the maximum of the minimum base sizes over all transitive representations of G , and suppose that there is an intransitive representation with minimum base size greater than $b_3^*(G)$. Now there exist at most $b_3(G)$ points in an orbit whose stabiliser acts trivially on that orbit, and hence is trivial (since the action on the orbit is faithful), contrary to assumption.

Now let $b_3^+(G)$ be the maximum of the minimum base sizes over all primitive representations of G , and suppose that there is a transitive but imprimitive representation with base size greater than $b_3^+(G)$. There are at most $b_3^+(G)$ maximal blocks whose stabiliser acts trivially on the block system containing them, and hence is trivial (since again the action is faithful), contrary to assumption. \square

This proposition does not hold for $b_2(G)$. For the group $G = \text{PSL}(2, 7) \cong \text{PSL}(3, 2)$, a minimal base in any transitive representation has size at most 3. However, in the action on the points and lines of the projective plane of order 2, there is a minimal base of size 4, consisting of two points and two lines such that each point lies on one of the lines and each line passes through one of the points.

3 Boolean semilattices

The subgroups of the group G form a lattice $L(G)$, with the operations $H \wedge K = H \cap K$ and $H \vee K = \langle H, K \rangle$. A *meet-semilattice* of $L(G)$ is a collection of subgroups containing G and closed under \wedge , while a *join-semilattice* is a collection of subgroups containing the trivial group 1 and closed under \vee .

The *Boolean lattice* $B(n)$ is the lattice of subsets of an n -set.

Proposition 3.1 *Let G be a finite group. Then $B(n)$ is embeddable as a meet-semilattice in $L(G)$ if and only if it is embeddable as a join-semilattice.*

Proof Suppose first that $B(n)$ is a join-semilattice of $L(G)$. Let $N = \{1, \dots, n\}$. Then, for every subset I of N , there is a subgroup H_I of G , and $H_{I \cup J} = \langle H_I, H_J \rangle$ for any two subsets I and J . Moreover, all these subgroups are distinct. In particular, $H_i \not\leq H_{N \setminus \{i\}}$ for all i (where H_i is shorthand for $H_{\{i\}}$); else

$$H_N = \langle H_i, H_{N \setminus \{i\}} \rangle = H_{N \setminus \{i\}},$$

contrary to assumption.

Let $K_i = H_{N \setminus \{i\}}$, and, for any $I \subseteq N$, put

$$K_I = \bigcap_{i \in I} K_i,$$

with the convention that $K_\emptyset = G$. We claim that all the subgroups K_I are distinct. Suppose that two of them are equal, say $K_I = K_J$. By interchanging I and J if necessary, we may assume that there exists $i \in I \setminus J$. But then $H_i \leq K_J$ while $H_i \not\leq K_I$, a contradiction.

Now it is clear that $K_I \cap K_J = K_{I \cup J}$, so that we have an embedding of $B(n)$ as meet-semilattice (where we have reversed the order-isomorphism to simplify the notation).

The reverse implication is proved by an almost identical argument. \square

Note that the conditions of the proposition are not equivalent to embeddability of $B(n)$ as a lattice. For example, if G is the quaternion group of order 8, then $B(2)$ is embeddable as both a meet-semilattice and a join-semilattice but not as a lattice.

Proposition 3.2 *Let G be a finite group.*

- (a) *The largest n such that $B(n)$ is embeddable as a join-semilattice of $L(G)$ is $\mu'(G)$.*
- (b) *The largest n such that $B(n)$ is embeddable as a meet-semilattice of $L(G)$ in such a way that the minimal element is a normal subgroup of G is $b_2(G)$.*

Proof (a) Let g_1, \dots, g_n be an independent set in G , where $n = \mu'(G)$. Let $N = \{1, \dots, n\}$. Then the subgroups $H_I = \langle g_i : i \in I \rangle$ for $I \subseteq N$ form a join-semilattice of $L(G)$ isomorphic to $B(n)$.

Conversely, suppose that we have a join-semilattice given by the subgroups H_I for $I \subseteq N$. Choose $g_i \notin H_{N \setminus \{i\}}$. Then clearly the elements g_1, \dots, g_n are independent.

(b) Let $(\alpha_1, \dots, \alpha_n)$ be a minimal base for G in some permutation representation. Let K_i be the stabiliser of α_i , and $K_I = \bigcap_{i \in I} K_i$ for $I \subseteq N = \{1, \dots, n\}$. Then the subgroups K_I form a meet-semilattice of $L(G)$. The subgroup K_N is the kernel of the permutation representation (by definition of a base), and so is a normal subgroup of G .

Conversely, suppose that we have a meet-semilattice given by the subgroups K_I for $I \subseteq N = \{1, \dots, n\}$, such that K_N is a normal subgroup of G ; the notation is chosen so that $K_I \cap K_J = K_{I \cup J}$. Now consider the permutation representation

on the union of the coset spaces of the subgroups K_i . Since the intersection of all these subgroups is normal, it is the kernel of the representation, and the points corresponding to the given subgroups form a base. It is minimal, since the intersection of fewer than n of the subgroups is not equal to K_N . \square

Corollary 3.3 $b_2(G) \leq \mu'(G)$ for any group G . \square

I do not know a group where the inequality is strict. Resolving this is equivalent to the following question. Let n be maximal such that $B(n)$ is embeddable as a meet-semilattice of $L(G)$. Is there an embedding of $B(n)$ for which the minimal element is normal?

From the above results, we conclude:

Corollary 3.4 $b_2(S_n) = b_3(S_n) = n - 1$.

Proof We have

$$n - 1 \leq b_3(S_n) \leq b_2(S_n) \leq \mu'(S_n) = n - 1,$$

where the first inequality holds because any base in the natural representation has size $n - 1$; the second is trivial; the third comes from the preceding Corollary; and the equality is Whiston's Theorem. \square

References

- [1] L. Babai, On the length of subgroup chains in the symmetric group, *Commun. Algebra* **14** (1986), 1729–1736.
- [2] P. J. Cameron and Ph. Cara, Independent generating sets and geometries for symmetric groups, *J. Algebra*, in press.
- [3] P. J. Cameron, R. Solomon and A. Turull, Chains of subgroups in symmetric groups, *J. Algebra* **127** (1989), 340–352.
- [4] P. Diaconis and L. Saloff-Coste, Walks on generating sets of groups, *Invent. Math.* **134** (1998), 251–299.
- [5] M. R. Jerrum, A compact representation for permutation groups, *J. Algorithms* **7** (1986), 60–78.
- [6] A. McIver and P. M. Neumann, Enumerating finite groups, *Quart. J. Math.* (2) **38** (1987), 473–488.
- [7] J. Whiston, Maximal independent generating sets of the symmetric group, *J. Algebra* **232** (2000), 255–268.