

# Exact solution of a model of a vesicle attached to a wall subject to mechanical deformation

A L Owczarek<sup>1</sup> and T Prellberg<sup>2</sup>

<sup>1</sup> Department of Mathematics and Statistics,  
The University of Melbourne, Parkville, Vic 3010, Australia.  
owczarek@unimelb.edu.au

<sup>2</sup> School of Mathematical Sciences  
Queen Mary University of London  
Mile End Road, London E1 4NS, UK  
t.prellberg@qmul.ac.uk

November 14, 2011

## Abstract

Area-weighted Dyck-paths are a two-dimensional model for vesicles attached to a wall. We model the mechanical response of a vesicle to a pulling force by extending this model.

We obtain an exact solution using two different approaches, leading to a  $q$ -deformation of an algebraic functional equation, and a  $q$ -deformation of a linear functional equation with a catalytic variable, respectively. While the non-deformed linear functional equation is solved by substitution of special values of the catalytic variable (the so-called “kernel method”), the  $q$ -deformed case is solved by iterative substitution of the catalytic variable.

Our model shows a non-trivial phase transition when a pulling force is applied. As soon as the area is weighted with non-unity weight, this transition vanishes.

## 1 Introduction

With the growing experimental ability to probe the behaviour of a single polymer under the influence of a stretching force [13, 12], there has been an effort by theoretical work to explore models that cover the possible scenarios encountered. For example, various directed path models of a polymer with one end, or both ends, grafted to a surface and being pulled at the end or in the middle, with the addition of an attractive contact interaction with the surface, have been studied in two and three dimensions [21, 22, 6, 1, 15]. Numerical work on self-avoiding walks models has also confirmed and expanded upon the predictions of these directed models [16, 17, 19, 20].

Another type of model, that of a lattice polygon that has been weighted by its length and area, can rather be used to model biological vesicles [18, 2, 9, 23, 3, 4]. For example, the effect of a restricted geometry such as a slit has recently been analysed [24]. Here, it is also natural to consider the effect of a pulling force of the surface of the vesicle. Hence, we study a

simple directed model of a two-dimensional vesicle that is being pulled away from a wall. The model is of interest physically as well as mathematically. It illustrates the solution of functional equations with catalytic variables under a  $q$ -deformation, when the solution gets expressed in terms of  $q$ -series rather than in terms of algebraic functions.

## 2 The Model




Figure 1: An area-weighted Dyck path with 12 steps and area of size 7, which is being pulled away from the surface at its right-most peak with force  $f$ . The height  $h$  of the right-most peak in this example is 2.

Dyck paths are directed walks on  $\mathbb{Z}^2$  starting at  $(0,0)$  and ending on the line  $y = 0$ , which have no vertices with negative  $y$ -coordinates, and which have steps in the  $(1,1)$  (up-step) and  $(1,-1)$  (down-step) directions [27]. Given a Dyck path  $\pi$ , we define the length  $n(\pi)$  to be half the number of its steps, and the area  $m(\pi)$  to be the sum of the starting heights of all up-steps in the path. This is sometimes also called the rank function of a Dyck path [8, 26], and is equivalent to the number of diamond plaquettes under the Dyck path. We further define the height  $h(\pi)$  of the right-most peak as the length of the final descent of the Dyck path, i.e. the maximal number of the last consecutive down-steps. An example of a Dyck path is given in Fig. 1.

Let  $\mathcal{G}$  be the set of Dyck paths, and define the generating function

$$G(t, q, \lambda) = \sum_{\pi \in \mathcal{G}} t^{n(\pi)} q^{m(\pi)} \lambda^{h(\pi)}. \quad (2.1)$$

The generating function for area-weighted Dyck paths [10, 7] has previously been studied. Dyck paths weighted by height of the right-most peak are related to Ballot paths, which have been previously studied as a model of pulled polymers [21]. Here we extend these works by combining and extending these models.

The key to obtaining a solution of this model is a combinatorial construction of Dyck paths which leads to a functional equation for the generating function  $G$ . We shall present two different

constructions leading to two structurally different functional equations.

### 3 $q$ -deformed Algebraic Functional Equation

To obtain the first functional equation, we decompose Dyck paths by considering the left-most down-step ending at height zero. This leads to a decomposition of a Dyck path in terms of two smaller Dyck paths. More precisely, except for the zero-step path, this decomposes a Dyck path uniquely into a Dyck path, followed by another Dyck path which is bracketed by up and down steps. This naturally leads to a functional equation which is quadratic in the generating functions,

$$G(t, q, \lambda) = 1 + \lambda t G(t, q, 1) G(qt, q, \lambda) . \quad (3.1)$$

Note that in the product  $\lambda t G(t, q, 1) G(qt, q, \lambda)$  the left Dyck path no longer contributes to the height of the right-most peak, leading to the factor  $G(t, q, 1)$ , and the bracketing of the right Dyck path by up and down steps increases the area proportional to the length of the path, as well as increases height and length by one, leading to the factor  $\lambda t G(qt, q, \lambda)$ .

#### 3.1 The Case $q = 1$

The functional equation (3.1) reads now

$$G(t, 1, \lambda) = 1 + \lambda t G(t, 1, 1) G(t, 1, \lambda) , \quad (3.2)$$

which after substitution of  $\lambda = 1$  can be readily solved to give

$$G(t, 1, 1) = \frac{1 - \sqrt{1 - 4t}}{2t} = \frac{2}{1 + \sqrt{1 - 4t}} , \quad (3.3)$$

which is the well-known generating function for Catalan numbers. Substitution of this result back into Eqn. (3.2) gives

$$G(t, 1, \lambda) = \frac{2}{2 - \lambda + \lambda \sqrt{1 - 4t}} . \quad (3.4)$$

For comparison with the result in Eqn. (3.11), note that a series expansion in  $\lambda$  gives

$$G(t, 1, \lambda) = \sum_{l=0}^{\infty} \lambda^l t^l G(t, 1, 1)^l . \quad (3.5)$$

#### 3.2 The case $q \neq 1$

Substituting  $\lambda = 1$  into the functional equation (3.1) gives

$$G(t, q, 1) = 1 + t G(t, q, 1) G(qt, q, 1) . \quad (3.6)$$

The Ansatz

$$G(t, q, 1) = \frac{H(qt, q)}{H(t, q)} \quad (3.7)$$

gives a linear functional equation for  $H(t, q)$ ,

$$H(qt, q) = H(t, q) + tH(q^2t, q) . \quad (3.8)$$

A solution is given by

$$H(t, q) = \sum_{k=0}^{\infty} \frac{(-t)^k q^{k(k-1)}}{(q; q)_k} , \quad (3.9)$$

where we have used the  $q$ -product notation

$$(t; q)_k = \prod_{j=0}^{k-1} (1 - tq^j) . \quad (3.10)$$

Here  $H(t, q) = {}_0\phi_1(-; 0; q, -t)$  is a basic hypergeometric series [11] also referred to as the  $q$ -Airy function [14].

Once  $G(t, q, 1)$  is known and substituted back into Eqn. (3.1), we can solve for  $G(t, q, \lambda)$  by iteration and obtain

$$G(t, q, \lambda) = \sum_{l=0}^{\infty} \lambda^l t^l q^{\binom{l}{2}} \frac{H(q^l t, q)}{H(t, q)} . \quad (3.11)$$

This iteration makes sense in a ring of formal power series. Alternatively, one can argue that the iteration converges for  $|q| < 1$ .

We note that the limit  $q \rightarrow 1$  can be obtained as follows. As

$$G(t, 1, 1) = \lim_{q \rightarrow 1} G(t, q, 1) = \lim_{q \rightarrow 1} \frac{H(qt, q)}{H(t, q)} , \quad (3.12)$$

we deduce that for fixed value of  $k$ ,

$$G(t, 1, 1) = \lim_{q \rightarrow 1} \frac{H(q^{k+1}t, q)}{H(q^k t, q)} \quad (3.13)$$

and thus

$$\lim_{q \rightarrow 1} \frac{H(q^l t, q)}{H(q, t)} = G(t, 1, 1)^l . \quad (3.14)$$

Therefore Eqn. (3.11) reduces to the result in Eqn. (3.5) as  $q \rightarrow 1$ .

## 4 $q$ -deformed Linear Functional Equation

To obtain the second functional equation, we consider how Dyck path configurations get changed by the insertion of a small pattern.

The right-most descent, together with the preceding up-step forms a *hook*. Inserting an up-step followed immediately by a down-step in the right-most descent creates a new Dyck path with a new hook. If the descent has length  $h$ , there are  $h + 1$  possibilities of insertion.

The functional equation corresponding to this construction is given by

$$G(t, q, \lambda) = 1 + \frac{\lambda t}{1 - q\lambda} G(t, q, 1) - \frac{q\lambda^2 t}{1 - q\lambda} G(t, q, q\lambda) . \quad (4.1)$$

The terms on the right-hand side of (4.1) can be most easily understood by considering

$$G(t, q, \lambda) = \sum_{h=0}^{\infty} \lambda^h G_h(t, q), \quad (4.2)$$

where  $G_h(t, q)$  are partial generating functions for fixed height  $h$ . The only Dyck path with a hook of height  $h = 0$  is a zero-step path. A Dyck path with a hook of height  $h \geq 1$  can only have been constructed from a Dyck path with a hook of height at least  $h - 1$ . By the construction, its area increases by an amount  $h - 1$ , and we thus find that for  $h \geq 1$

$$G_h(t, q) = q^{h-1} \sum_{k=h-1}^{\infty} G_k(t, q). \quad (4.3)$$

Summing over the partial generating functions leads to the functional equation (4.1).

#### 4.1 The Case $q = 1$

The functional equation (4.1) now reads

$$G(t, 1, \lambda) = 1 + \frac{\lambda t}{1 - \lambda} G(t, 1, 1) - \frac{\lambda^2 t}{1 - \lambda} G(t, 1, \lambda). \quad (4.4)$$

Here, substitution of  $\lambda = 1$  is no longer useful, and one solves the functional equation using the kernel method [25]. Writing Eqn. (4.1) as

$$K(t, \lambda) G(t, 1, \lambda) = 1 + \frac{\lambda t}{1 - \lambda} G(t, 1, 1), \quad (4.5)$$

where the kernel  $K(t, \lambda)$  is given as

$$K(t, \lambda) = 1 + \frac{\lambda^2 t}{1 - \lambda}, \quad (4.6)$$

we observe that the left-hand side of Eqn. (4.5) vanishes if  $\lambda = \lambda_0$  is chosen such that  $K(t, \lambda_0)$  vanishes (provided  $G(t, 1, \lambda_0)$  does not diverge). This then implies that

$$G(t, 1, 1) = \frac{\lambda_0 - 1}{\lambda_0 t}, \quad (4.7)$$

and substituting this result back into Eqn. (4.5) leads to an expression for  $G(t, 1, \lambda)$ . The appropriate root of the kernel is given by

$$\lambda_0 = \frac{1 - \sqrt{1 - 4t}}{2t}, \quad (4.8)$$

Using this value, we recover the expressions (3.3) for  $G(t, 1, 1)$  and (3.4) for  $G(t, 1, \lambda)$  given above.

## 4.2 The Case $q \neq 1$

If  $q \neq 1$ , we can solve the functional equation (4.1) by iterative substitution. Substituting  $\lambda q^k$  for  $\lambda$  in (4.1), we find

$$G(t, q, q^k \lambda) = 1 + \frac{q^k \lambda t}{1 - q^{k+1} \lambda} G(t, q, 1) - \frac{q^{2k+1} \lambda^2 t}{1 - q^{k+1} \lambda} G(t, q, q^{k+1} \lambda). \quad (4.9)$$

We can therefore express  $G(t, q, q^k \lambda)$  in terms of  $G(t, q, q^{k+1} \lambda)$  and use this to iterate. As a result, we find

$$G(t, q, \lambda) = \sum_{k=0}^{\infty} \frac{(-\lambda^2 t)^k q^{k^2}}{(q\lambda; q)_k} \left( 1 + \frac{q^k \lambda t}{1 - q^{k+1} \lambda} G(t, q, 1) \right). \quad (4.10)$$

Defining

$$I(t, q, y) = \sum_{k=0}^{\infty} \frac{(-t)^k q^{k(k-1)}}{(y; q)_k}, \quad (4.11)$$

this simplifies to

$$G(t, q, \lambda) = I(q\lambda^2 t, q, q\lambda) - \frac{1}{\lambda} [I(\lambda^2 t, q, q\lambda) - 1] G(t, q, 1). \quad (4.12)$$

Note that  $I(t, q, y) = {}_1\phi_2(q; 0, y; q, -t)$  is also expressible as a basic hypergeometric function. Moreover, it is a deformation of  $H(t, q)$ , in that  $I(t, q, q) = H(t, q)$ .

Substituting  $\lambda = 1$  in Eqn. (4.12) gives now immediately

$$G(t, q, 1) = \frac{H(qt, q)}{H(t, q)}, \quad (4.13)$$

a result already obtained from the  $q$ -algebraic functional equation (3.6) above. Substituting this result back into Eqn. (4.12) leads to the main result of this section, namely

$$G(t, q, \lambda) = I(q\lambda^2 t, q, q\lambda) - \frac{1}{\lambda} [I(\lambda^2 t, q, q\lambda) - 1] \frac{H(qt, q)}{H(t, q)}. \quad (4.14)$$

Note the rather different structure of the equivalent solutions (3.11) and (4.14). The former is in terms of an infinite sum, which however readily allows for extraction of the partial generating function for fixed height  $h$ . The latter is more compact as it is written in terms of a finite number of special functions. We note that the equivalence of both solutions implies the identity

$$\sum_{l=0}^{\infty} \lambda^l t^l q^{\binom{l}{2}} H(q^l t, q) = I(q\lambda^2 t, q, q\lambda) H(t, q) - \frac{1}{\lambda} [I(\lambda^2 t, q, q\lambda) - 1] H(qt, q). \quad (4.15)$$

## 5 Analysis of behaviour

Let us begin by considering the case of  $q = 1$ , that is no extra area weighting or, equivalently, osmotic pressure. The first important point to make is that previous models of pulling on the end of a polymer without any other interaction with a surface give the result that as soon as a force is applied the end of the polymer changes its scaling behaviour from random-walk scaling

$$\langle h \rangle \sim n^{1/2} \quad (5.1)$$




Figure 2: A plot of the radius of convergence  $t_c(\lambda) \equiv t_c(\lambda, 1)$  in the polymer length variable  $t$  of the generating function when  $q = 1$  as a function of the force variable  $\lambda$ . Note that there is a change of behaviour at  $\lambda = 2$  with constant  $t_c(\lambda) = 1/4$  for  $\lambda \leq 2$ .

for  $f = 0$  ( $\lambda = 1$ ) to ballistic scaling

$$\langle h \rangle \sim n \quad (5.2)$$

for  $f > 0$  ( $\lambda > 1$ ). This is reflected in the fact that the radius of convergence of the associated generating function is constant for  $\lambda \leq 1$  and decreases continuously for  $\lambda > 1$ . As a model of pulling on a vesicle attached to a surface, our model has the feature that the pulling occurs on the final “peak”, the location of which along the surface can change with the pulling itself. As such, there is a non-trivial critical point as a function of  $\lambda$  in our model, with a critical value

$$\lambda = \lambda_c \equiv 2. \quad (5.3)$$

A plot of the radius of convergence of the generating function (3.4) is shown in Fig. 2.

Note that the reduced free energy  $\kappa(\lambda, q)$  of the surface is simply related to the radius of convergence  $t_c(\lambda, q)$  in the usual way via

$$\kappa(\lambda, q) = -\log t_c(\lambda, q). \quad (5.4)$$

For  $q < 1$  the generating function (3.11) can be seen to be convergent except where  $H(t, q) = 0$ . Crucially this is independent of  $\lambda$ ! This implies that for  $q < 1$  there is no phase transition as a function of  $\lambda$ , since  $\kappa(\lambda, q)$  is independent of  $\lambda$  for  $q < 1$ . Physically this means that the osmotic pressure always dominates the pulling force for  $q < 1$ :  $\langle h \rangle$  will be bounded for all  $\lambda$ .

Interestingly, if one considers the limit  $q \rightarrow 1$  for  $\lambda > 2$  one is left with the problem that the free energy “jumps” at  $q = 1$ . The resolution of this apparent problem is that for  $q > 1$  one needs to consider the free energy per unit area rather than the free energy per unit length of the surface. The limiting free energy per unit area is a continuous function of  $q$  and  $\lambda$  for all  $\lambda$ , being equal to zero for  $q \leq 1$ .

## 6 Conclusion

We have provided the exact solution of a simple model of a vesicle attached to a surface subject to a pulling force on the vesicle. We have used two methods of solution which illuminate the generalisation of the well-known “kernel” method for the solution of linear functional equations using a “catalytic” variable and result in a non-trivial identity. The solution is expressed in terms of well-studied  $q$ -series and can be analysed simply. Our model has a non-trivial phase transition when no osmotic pressure is applied.

## Acknowledgements

Financial support from the Australian Research Council via its support for the Centre of Excellence for Mathematics and Statistics of Complex Systems is gratefully acknowledged by the authors. A L Owczarek thanks the School of Mathematical Sciences, Queen Mary, University of London for hospitality.

## References

- [1] J. Alvarez and S. G. Whittington, *Force-extension relations for adsorbing polymers subject to a force*, J. Stat. Mech. (2009) P04016.
- [2] R. Brak and A. J. Guttmann, *Exact solution of a staircase and row-convex polygon perimeter and area generating function*, J. Phys. A. **23** (1990) 4581.
- [3] R. Brak, A. L. Owczarek, and T. Prellberg, *Exact Scaling Behaviour of Partially Convex Vesicles*, J. Stat. Phys. **76** (1994) 1101.
- [4] R. Brak, J. M. Essam, and A. L. Owczarek, *New results for directed vesicles and chains near an attractive wall*, J. Stat. Phys. **93** (1998) 155.
- [5] R. Brak, A. L. Owczarek, A. Rechnitzer, and S. Whittington, *A directed walk model of a long chain polymer in a slit with attractive walls*, J. Phys. A **38** (2005) 4309.
- [6] R. Brak, P. Dyke, J. Lee, A. L. Owczarek, T. Prellberg, A. Rechnitzer, and S. Whittington, *A self-interacting partially directed walks subject to a force*, J. Phys. A **42** (2009) 085001.
- [7] P. Duchon, *On the Enumeration and Generation of Generalized Dyck Words*, Discrete Mathematics **225** (2000) 121.

- [8] L. Ferrari and R. Pinzani, *Lattices of lattices paths*, J. Statist. Plann. Inference **135** (2005) 7792.
- [9] M. E. Fisher, A. J. Guttmann, and S. Whittington, J. Phys. A **24** (1991) 3095.
- [10] P. Flajolet, *Combinatorial aspects of continued fractions*, Discrete Mathematics **41** (1980) 125.
- [11] G. Gasper and M. Rahman, *Basic Hypergeometric Series*, volume 96 of *Encyclopedia of Mathematics and its Applications*, Cambridge University Press, Cambridge, 2004.
- [12] H. Hauchke, M. J. Miles, and V. Koutsos, *Conformation of a single polyacrylamide molecule adsorbed onto a mica surface studied with atomic force microscopy*, Macromolecules **37** (2004) 3799.
- [13] B. J. Haupt, J. Ennis, and E. M. Sevick, *The detachment of a polymer chain from a weakly adsorbing surface using an AFM tip*, Langmuir **15** (1999) 3886.
- [14] M. E. H. Ismail *Asymptotics of  $q$ -orthogonal polynomials and a  $q$ -Airy function* Int. Math. Res. Notices **2005** (2005) 1063.
- [15] E. J. Janse van Rensburg, *Pulled directed lattice paths*, J. Phys. A **43** (2010) 215001.
- [16] J. Krawczyk, T. Prellberg, A. L. Owczarek, and A. Rechnitzer, *Stretching of a chain polymer adsorbed at a surface*, J. Stat. Mech. (2004) P10004.
- [17] J. Krawczyk, A. L. Owczarek, T. Prellberg, and A. Rechnitzer, *Pulling absorbing and collapsing polymers from a surface*, J. Stat. Mech. (2005) P05008.
- [18] S. Leibler, R. R. P. Singh, and M. E. Fisher, *Thermodynamic behavior of two-dimensional vesicles*, Phys. Rev. Lett. **59** (1987) 1989.
- [19] P. K. Mishra, S. Kumar, and Y. Singh, *Force-induced desorption of a linear polymer chain adsorbed on an attractive surface*, Europhys. Lett. **69** (2005) 102.
- [20] E. Orlandini and M. C. Tesi, *Modelling the adsorption of a polymer subject to an elongational force by directed walk models*, J. Math. Chem. **45** (2009) 72.
- [21] E. Orlandini and S. G. Whittington, *Pulling a polymer at an interface: directed walk models*, J. Phys. A **37** (2004) 5305.
- [22] E. Orlandini, M. C. Tesi, and S. G. Whittington, *Adsorption of a directed polymer subject to an elongational force*, J. Phys. A **37** (2004) 1535.
- [23] A. L. Owczarek and T. Prellberg, *Exact Solution of the Discrete (1+1)-Dimensional SOS Model with Field and Surface Interactions*, J. Stat. Phys. **70** (1993) 1175.

- [24] A. L. Owczarek and T. Prellberg, *A simple model of a vesicle drop in a confined geometry*, J. Stat. Mech. (2010) P08015.
- [25] H. Prodinger, *The Kernel Method: A Collection of Examples*, Sém. Lothar. Combin. **50** (2004) B50f.
- [26] A. Sapounakis, I. Tasoulas, and P. Tsikouras, *On the Dominance Partial Ordering of Dyck Paths*, Journal of Integer Sequences **9** (2006) 06.2.5.
- [27] R. P. Stanley, *Enumerative Combinatorics, Volume 2*, volume 62 of *Cambridge Studies in Advanced Mathematics*, Cambridge University Press, Cambridge, 1999.